

USSN: 10/643,681
Amdt. Dated June 7, 2004
Page 3 of 10

Amendments to the Claims:

This listing of claims will replace all prior versions and listings of claims in the application:

Listing of Claims:

Claims 1-23 cancelled.

24. (New) A method of reducing or moderating a postprandial rise in plasma glucose in a mammal comprising administering to said mammal an amylin or an amylin agonist.

25. (New) The method of claim 24 wherein the amylin agonist is an amylin agonist analogue having the following amino acid sequence:

¹A₁-X-Asn-Thr-⁵Ala-Thr-Y-Ala-Thr¹⁰Gln-Arg-Leu-B₁-Asn-¹⁵Phe-Leu-C₁-D₁-E₁-²⁰F₁-G₁-Asn-H₁-Gly-²⁵Pro-I₁-Leu-Pro-J₁-³⁰Thr-K₁-Val-Gly-Ser-³⁵Asn-Thr-Tyr-Z

wherein

A₁ is Lys, Ala Ser or Hydrogen;

B₁ is Ala, Ser or Thr;

C₁ is Val, Leu or Ile;

D₁ is His or Arg;

E₁ is Ser or Thr;

F₁ is Ser, Thr, Gln or Asn;

G₁ is Asn, Gln or His;

H₁ is Phe, Leu or Tyr;

I₁ is Ile, Val, Ala or Leu

J₁ is Ser, Pro or Thr;

K₁ is Asn, Asp or Gln;

X and Y are independently selected residues having side chains which are chemically bonded to each other to form an intramolecular linkage, wherein said intramolecular linkage comprises a disulfide bond, a lactam or a thioether linkage; and Z is an amino, alkylamino, dialkylamino, cycloalkylamino, arylamino, aralkylamino, alkyloxy, aryloxy or aralkyloxy; and provided that when A₁ is Lys, B₁ is Ala, C₁ is Val, D₁ is Arg, E₁ is Ser, F₁ is Ser, G₁ is Asn, H₁ is Leu, I₁ is Val, J₁ is Pro, and K₁ is Asn; then one or more A₁ to K₁ is a D-amino acid and Z is selected from the group consisting of Alkylamino, dialkylamino, cycloalkylamino, arylamino, aralkylamino, alkyloxy, aryloxy or aralkyloxy.

USSN: 10/643,681
 Amdt. Dated June 7, 2004
 Page 4 of 10

26. (New) The method of claim 24 wherein the amylin agonist is an amylin agonist analogue having the following amino acid sequence:

¹A₁-X-Asn-Thr-⁵Ala-Thr-Y-Ala-Thr-¹⁰Gln-Arg-Leu-B₁-Asn-¹⁵Phe-Leu-C₁-D₁-E₁-²⁰F₁-
 G₁-Asn-H₁-Gly-²⁵Pro-I₁-Leu-J₁-Pro-³⁰Thr-K₁-Val-Gly-Ser-³⁵Asn-Thr-Tyr-Z

wherein

A₁ is Lys, Ala, Ser or hydrogen;

B₁ is Ala, Ser or Thr;

C₁ is Val, Leu or Ile;

D₁ is His or Arg;

E₁ is Ser or Thr;

F₁ is Ser, Thr, Gln or Asn;

G₁ is Asn, Gln or His;

H₁ is Phe, Leu or Tyr;

I₁ is Ile, Val, Ala or Leu;

J₁ is Ser, Pro, Leu, Ile or Thr;

K₁ is Asn, Asp or Gln;

X and Y are independently selected residues having side chains which are chemically bonded to each other to form an intramolecular linkage, wherein said intramolecular linkage comprises a disulfide bond, a lactam or a thioether linkage; and Z is amino, alkylamino, dialkylamino, cycloalkylamino, arylamino, aralkylamino, alkyloxy, aryloxy or aralkyloxy; and provided that when

(a) A₁ is Lys, B₁ is Ala, C₁ is Val, D₁ is Arg, E₁ is Ser, F₁ is Ser, G₁ is Asn, H₁ is Leu, I₁ is Val, J₁ is Pro and K₁ is Asn; or

(b) A₁ is Lys, B₁ is Ala, C₁ is Val, D₁ is His, E₁ is Ser, F₁ is Asn, G₁ is Asn, H₁ is Leu, I₁ is Val, J₁ is Ser and K₁ is Asn;

then one or more of A₁ to K₁ is a D-amino acid and Z is selected from the group consisting of alkylamino, dialkylamino, cycloalkylamino, arylamino, aralkylamino, alkyloxy, aryloxy or aralkyloxy.

27. (New) The method of claim 24 wherein the amylin agonist is an amylin agonist analogue having the following amino acid sequence:

¹A₁-X-Asn-Thr-⁵Ala-Thr-Y-Ala-Thr-¹⁰Gln-Arg-Leu-B₁-Asn-¹⁵Phe-Leu-C₁-D₁-E₁-²⁰F₁-
 G₁-Asn-H₁-Gly-²⁵I₁-J₁-Leu-Pro-Pro-³⁰Thr-K₁-Val-Gly-Ser-³⁵Asn-Thr-Tyr-Z

wherein

USSN: 10/643,681
Amdt. Dated June 7, 2004
Page 5 of 10

A₁ is Lys, Ala, Ser or hydrogen;

B₁ is Ala, Ser or Thr;

C₁ is Val, Leu or Ile;

D₁ is His or Arg;

E₁ is Ser or Thr;

F₁ is Ser, Thr, Gln or Asn;

G₁ is Asn, Gln or His;

H₁ is Phe, Leu or Tyr;

I₁ is Ala or Pro;

J₁ is Ile, Val, Ala or Leu;

K₁ is Asn, Asp or Gln;

X and Y are independently selected residues having side chains which are chemically bonded to each other to form an intramolecular linkage, wherein said intramolecular linkage comprises a disulfide bond, a lactam or a thioether linkage; and Z is amino, alkylamino dialkylamino, cycloalkylamino, arylamino, aralkylamino, alkylloxy, aryloxy or aralkylloxy; and provided that when A₁ is Lys, B₁ is Ala, C₁ is Val, D₁ is Arg, E₁ is Ser, F₁ is Ser, G₁ is Asn H₁ is Leu, I₁ is Pro, J₁ is Val and K₁ is Asn; then one or more of A₁ to K₁ is a D-amino acid and Z is selected from the group consisting of alkylamino, dialkylamino, cycloalkylamino, arylamino, aralkylamino, alkylloxy, aryloxy or aralkylloxy.

28. (New) The method of claim 24 wherein the amylin agonist is an amylin agonist analogue having the following amino acid sequence:

¹A₁-X-Asn-Thr-⁵Ala-Thr-Y-Ala-Thr-¹⁰Gln-Arg-Leu-B₁-Asn-¹⁵Phe-Leu-C₁-D₁-E₁-²⁰F₁-G₁-Asn-H₁-Gly-²⁵Pro-I₁-Leu-Pro-Pro-³⁰Thr-J₁-Val-Gly-Ser-³⁵Asn-Thr-Tyr-Z

wherein

A₁ is Lys, Ala, Ser or hydrogen;

B₁ is Ala, Ser or Thr;

C₁ is Val, Leu or Ile;

D₁ is His or Arg;

E₁ is Ser or Thr;

F₁ is Ser, Thr, Gln or Asn;

G₁ is Asn, Gln or His;

H₁ is Phe, Leu or Tyr;

USSN: 10/643,681
 Amdt. Dated June 7, 2004
 Page 6 of 10

I₁ is Ile, Val, Ala or Leu;

J₁ is Asn, Asp or Gln;

X and Y are independently selected residues having side chains which are chemically bonded to each other to form an intramolecular linkage wherein said intramolecular linkage comprises a disulfide bond, a lactam or a thioether linkage; and Z is amino, alkylamino, dialkylamino, cycloalkylamino, arylamino, aralkylamino, alkyloxy, aryloxy or aralkyloxy; and

provided that when A₁ is Lys, B₁ is Ala, C₁ is Val, D₁ is Arg, E₁ is Ser, F₁ is Ser, G₁ is Asn, H₁ is Leu, I₁ is Val and J₁ is Asn; then one or more of A₁ to J₁ is a D-amino acid and Z is selected from the group consisting of alkylamino, dialkylamino, cycloalkylamino, arylamino, aralkylamino, alkyloxy, aryloxy or aralkyloxy.

29. (New) The method of claim 24 wherein said amylin agonist is any one of ¹⁸Arg^{25,28}Pro-h-amylin, des-¹Lys¹⁸Arg^{25,28}Pro-h-amylin, ^{25,28,29}Pro-h-amylin, des-¹Lys^{25,28,29}Pro-h-amylin, ¹⁸Arg^{25,28,29}Pro-h-amylin, des-¹Lys¹⁸Arg^{25,28,29}Pro-h-amylin, ²⁵Pro²⁶Val^{28,29}Pro-h-amylin, or des-¹Lys²⁵Pro²⁶Val^{28,29}Pro-h-amylin.

30. (New) The method of claim 24 wherein the amylin agonist is ^{25,28,29}Pro-h-amylin.

31. (New) A method of treating ingestion of a toxin in a mammal comprising administering to said mammal an amylin or an amylin agonist and aspirating the toxin out of a stomach of the mammal.

32. (New) The method of claim 31 wherein the amylin agonist is an amylin agonist analogue having the following amino acid sequence:

¹A₁-X-Asn-Thr-⁵Ala-Thr-Y-Ala-Thr¹⁰Gln-Arg-Leu-B₁-Asn-¹⁵Phe-Leu-C₁-D₁-E₁-²⁰F₁-G₁-Asn-H₁-Gly-²⁵Pro-I₁-Leu-Pro-J₁-³⁰Thr-K₁-Val-Gly-Ser-³⁵Asn-Thr-Tyr-Z

wherein

A₁ is Lys, Ala Ser or Hydrogen;

B₁ is Ala, Ser or Thr;

C₁ is Val, Leu or Ile;

D₁ is His or Arg;

E₁ is Ser or Thr;

F₁ is Ser, Thr, Gln or Asn;

G₁ is Asn, Gln or His;

USSN: 10/643,681
Amdt. Dated June 7, 2004
Page 7 of 10

H₁ is Phe, Leu or Tyr;

I₁ is Ile, Val, Ala or Leu

J₁ is Ser, Pro or Thr;

K₁ is Asn, Asp or Gln;

X and Y are independently selected residues having side chains which are chemically bonded to each other to form an intramolecular linkage, wherein said intramolecular linkage comprises a disulfide bond, a lactam or a thioether linkage; and Z is an amino, alkylamino, dialkylamino, cycloalkylamino, arylamino, aralkylamino, alkyloxy, aryloxy or aralkyloxy; and provided that when A₁ is Lys, B₁ is Ala, C₁ is Val, D₁ is Arg, E₁ is Ser, F₁ is Ser, G₁ is Asn, H₁ is Leu, I₁ is Val, J₁ is Pro, and K₁ is Asn; then one or more A₁ to K₁ is a D-amino acid and Z is selected from the group consisting of Alkylamino, dialkylamino, cycloalkylamino, arylamino, aralkylamino, alkyloxy, aryloxy or aralkyloxy.

33. (New) The method of claim 31 wherein the amylin agonist is an amylin agonist analogue having the following amino acid sequence:

¹A₁-X-Asn-Thr-⁵Ala-Thr-Y-Ala-Thr-¹⁰Gln-Arg-Leu-B₁-Asn-¹⁵Phe-Leu-C₁-D₁-E₁-²⁰F₁-
G₁-Asn-H₁-Gly-²⁵Pro-I₁-Leu-J₁-Pro-³⁰Thr-K₁-Val-Gly-Ser-³⁵Asn-Thr-Tyr-Z

wherein

A₁ is Lys, Ala, Ser or hydrogen;

B₁ is Ala, Ser or Thr;

C₁ is Val, Leu or Ile;

D₁ is His or Arg;

E₁ is Ser or Thr;

F₁ is Ser, Thr, Gln or Asn;

G₁ is Asn, Gln or His;

H₁ is Phe, Leu or Tyr;

I₁ is Ile, Val, Ala or Leu;

J₁ is Ser, Pro, Leu, Ile or Thr;

K₁ is Asn, Asp or Gln;

X and Y are independently selected residues having side chains which are chemically bonded to each other to form an intramolecular linkage, wherein said intramolecular linkage comprises a disulfide bond, a lactam or a thioether linkage; and Z is amino, alkylamino, dialkylamino, cycloalkylamino, arylamino, aralkylamino, alkyloxy, aryloxy or aralkyloxy; and provided that when

(c) A₁ is Lys, B₁ is Ala, C₁ is Val, D₁ is Arg, E₁ is Ser, F₁ is Ser, G₁ is Asn, H₁ is Leu, I₁ is Val, J₁ is Pro and K₁ is Asn; or

USSN: 10/643,681
Amdt. Dated June 7, 2004
Page 8 of 10

(d) A₁ is Lys, B₁ is Ala, C₁ is Val, D₁ is His, E₁ is Ser, F₁ is Asn, G₁ is Asn, H₁ is Leu, I₁ is Val, J₁ is Ser and K₁ is Asn;

then one or more of A₁ to K₁ is a D-amino acid and Z is selected from the group consisting of alkylamino, dialkylamino, cycloalkylamino, arylamino, aralkylamino, alkyloxy, aryloxy or aralkyloxy.

34. (New) The method of claim 31 wherein the amylin agonist is an amylin agonist analogue having the following amino acid sequence:

¹A₁-X-Asn-Thr-⁵Ala-Thr-Y-Ala-Thr-¹⁰Gln-Arg-Leu-B₁-Asn-¹⁵Phe-Leu-C₁-D₁-E₁-²⁰F₁-G₁-Asn-H₁-Gly-²⁵I₁-J₁-Leu-Pro-Pro-³⁰Thr-K₁-Val-Gly-Ser-³⁵Asn-Thr-Tyr-Z

wherein

A₁ is Lys, Ala, Ser or hydrogen;

B₁ is Ala, Ser or Thr;

C₁ is Val, Leu or Ile;

D₁ is His or Arg;

E₁ is Ser or Thr;

F₁ is Ser, Thr, Gln or Asn;

G₁ is Asn, Gln or His;

H₁ is Phe, Leu or Tyr;

I₁ is Ala or Pro;

J₁ is Ile, Val, Ala or Leu;

K₁ is Asn, Asp or Gln;

X and Y are independently selected residues having side chains which are chemically bonded to each other to form an intramolecular linkage, wherein said intramolecular linkage comprises a disulfide bond, a lactam or a thioether linkage; and Z is amino, alkylamino, dialkylamino, cycloalkylamino, arylamino, aralkylamino, alkyloxy, aryloxy or aralkyloxy; and provided that when A₁ is Lys, B₁ is Ala, C₁ is Val, D₁ is Arg, E₁ is Ser, F₁ is Ser, G₁ is Asn, H₁ is Leu, I₁ is Pro, J₁ is Val and K₁ is Asn; then one or more of A₁ to K₁ is a D-amino acid and Z is selected from the group consisting of alkylamino, dialkylamino, cycloalkylamino, arylamino, aralkylamino, alkyloxy, aryloxy or aralkyloxy.

35. (New) The method of claim 31 wherein the amylin agonist is an amylin agonist analogue having the following amino acid sequence:

¹A₁-X-Asn-Thr-⁵Ala-Thr-Y-Ala-Thr-¹⁰Gln-Arg-Leu-B₁-Asn-¹⁵Phe-Leu-C₁-D₁-E₁-²⁰F₁-G₁-Asn-H₁-Gly-²⁵Pro-I₁-Leu-Pro-Pro-³⁰Thr-J₁-Val-Gly-Ser-³⁵Asn-Thr-Tyr-Z

USSN: 10/643,681
Amdt. Dated June 7, 2004
Page 9 of 10

wherein

A₁ is Lys, Ala, Ser or hydrogen;

B₁ is Ala, Ser or Thr;

C₁ is Val, Leu or Ile;

D₁ is His or Arg;

E₁ is Ser or Thr;

F₁ is Ser, Thr, Gln or Asn;

G₁ is Asn, Gln or His;

H₁ is Phe, Leu or Tyr;

I₁ is Ile, Val, Ala or Leu;

J₁ is Asn, Asp or Gln;

X and Y are independently selected residues having side chains which are chemically bonded to each other to form an intramolecular linkage wherein said intramolecular linkage comprises a disulfide bond, a lactam or a thioether linkage; and Z is amino, alkylamino, dialkylamino, cycloalkylamino, arylamino, aralkylamino, alkyloxy, aryloxy or aralkyloxy; and

provided that when A₁ is Lys, B₁ is Ala, C₁ is Val, D₁ is Arg, E₁ is Ser, F₁ is Ser, G₁ is Asn, H₁ is Leu, I₁ is Val and J₁ is Asn; then one or more of A₁ to J₁ is a D-amino acid and Z is selected from the group consisting of alkylamino, dialkylamino, cycloalkylamino, arylamino, aralkylamino, alkyloxy, aryloxy or aralkyloxy.

36. (New) The method of claim 31 wherein said amylin agonist is any one of ¹⁸Arg^{25,28}Pro-h-amylin, des-¹Lys¹⁸Arg^{25,28}Pro-h-amylin, ^{25,28,29}Pro-h-amylin, des-¹Lys^{25,28,29}Pro-h-amylin, ¹⁸Arg^{25,28,29}Pro-h-amylin, des-¹Lys¹⁸Arg^{25,28,29}Pro-h-amylin, ²⁵Pro²⁶Val^{28,29}Pro-h-amylin, or des-¹Lys²⁵Pro²⁶Val^{28,29}Pro-h-amylin.

37. (New) The method of claim 31 wherein the amylin agonist is ^{25,28,29}Pro-h-amylin.